

Deformations of Flows from Type IIB Supergravity

Changhyun Ahn ^{*,†} and Justin F. Vázquez-Poritz ^{*,‡}

^{*} *School of Natural Sciences, Institute for Advanced Study,
Einstein Drive, Princeton NJ 08540, USA*

[†] *Department of Physics, Kyungpook National University,
Taegu 702-701, Korea*

[‡] *Department of Physics, University of Cincinnati,
Cincinnati OH 45221-001, USA*

ahn@knu.ac.kr, jporitz@physics.uc.edu

Abstract

We consider supersymmetric $SL(3, R)$ deformations of various type IIB supergravity backgrounds which exhibit flows away from an asymptotically locally $AdS_5 \times S^5$ fixed point. This includes the gravity dual of the Coulomb branch of $\mathcal{N} = 1$ super Yang Mills theory, for which the deformed superpotential is known. We also consider the gravity duals of field theories which live on various curved backgrounds, such as $Minkowski_2 \times H^2$, $AdS_3 \times S^1$ and $\mathbf{R} \times S^3$. Some of the deformed theories flow from a four-dimensional $\mathcal{N} = 1$ superconformal UV fixed point to a two-dimensional $(2, 2)$ superconformal IR fixed point. We study nonsupersymmetric generalizations of the deformations of the above Coulomb branch flows.

1 Introduction

Compactification of type IIB supergravity on a two-torus has an $SL(3, R)$ symmetry which can be used for generating new solutions. This has recently been applied to finding the type IIB supergravity background which, via the AdS/CFT correspondence [1, 2, 3], corresponds to marginal deformations of $\mathcal{N} = 4$ super Yang-Mills theory [4]. The undeformed theory has an isometry group which includes $U(1) \times U(1)$. The deformation on the gravity side can be matched to an exactly marginal operator in the field theory, providing a holographic test of the methods of Leigh and Strassler [5].

The solution-generating technique can be outlined as follows. First, T-dualize along one of the $U(1)$ directions to type IIA theory. Lifting the solution to eleven dimensions provides a third direction which is associated with a $U(1)$ symmetry. One can now apply an $SL(3, R)$ rotation along these $U(1)^3$ directions. Dimensionally reducing and T-dualizing along shifted directions yields a new type IIB solution. This procedure can be applied to any solution that has an isometry group which contains $U(1) \times U(1)$. If in addition to this symmetry there is a $U(1)$ R-symmetry, then the deformed solution preserves $\mathcal{N} = 1$ supersymmetry.

This method has also been applied to the marginal deformations of the conifold superconformal field theory [4], as well as the superconformal theories associated to the $Y^{p,q}$ [4] and $L^{p,q,r}$ [6] manifolds¹. Furthermore, the gravity duals of non-conformal theories, such as those which exhibit renormalization group flows, can also be deformed in this manner. In particular, deformations of the Klebanov-Strassler solution [11] and the 5-brane wrapped on a 2-sphere [12] have been considered in [4] and [13], respectively². These solutions are useful for understanding the gravity dual of confining gauge theories.

Marginal deformations of eleven-dimensional supergravity solutions with $U(1)^3$ symmetry have also been studied. The solution-generating technique is analogous to the above type IIB cases. One begins with a solution that has $U(1)^3$ non-R symmetry. Then one reduces to ten dimensions and T-dualizes to type IIB theory, where one can perform an $SL(2, R)$ transformation with the remaining two $U(1)$ directions and then dualize back to eleven dimensions along the shifted directions³. This has been done for initial solutions with geometries of the form $AdS_4 \times X^7$, which correspond to three-dimensional superconformal field theories. The

¹The Sasaki-Einstein spaces $Y^{p,q}$ and $L^{p,q,r}$ were found in [7, 8] and [9, 10], respectively.

²For the latter case, deformations which incorporate one internal direction and one worldvolume direction were also considered. These correspond to dipole deformations [13].

³In fact, there is also a direction associated with the $U(1)$ R-symmetry. This can be used to perform a multiple chain of dualities and $SL(3, R)$ transformations resulting in four deformation parameters. However, the deformed solution would be nonsupersymmetric. One could consider an eleven-dimensional solution with a large enough isometry group such that multiple-parameter deformations preserve supersymmetry.

case in which X^7 is a 7-sphere was considered in [4], for which the deformed gauge theory has $\mathcal{N} = 2$ supersymmetry and the exactly marginal deformation operator was identified. The deformed field theory also has $\mathcal{N} = 2$ supersymmetry if X^7 is one of a countably infinite number of Sasaki-Einstein spaces⁴ [6] and has $\mathcal{N} = 1$ supersymmetry if X^7 is weak G_2 or tri-Sasakian [16].

In this paper, we consider deformations of type IIB solutions which are asymptotically locally $AdS_5 \times S^5$. This implies that the UV limit of the dual gauge theory is the Leigh-Strassler deformation of $\mathcal{N} = 4$ super Yang-Mills theory. There is a renormalization group flow away from this conformal fixed point. We consider type IIB solutions which are related to solutions of the $U(1)^3$ truncation of five-dimensional gauged supergravity by using the dimensional reduction ansatz given in [17]. This implies that, when lifted to ten-dimensional type IIB theory, the isometries includes a $U(1)^2$ symmetry with which to deform the solution while maintaining a $U(1)$ R-symmetry. That is, at every point along the flow there is locally an S^5 which can be written as a constant $U(1)$ bundle over $\mathbb{CP}^2$. It is this $U(1)$ direction which corresponds to the $U(1)$ R-symmetry after the deformations. Non-vanishing $U(1)$ gauge fields in the five-dimensional solutions correspond to nonzero R -symmetry currents in the dual gauge theory. Although we have not explicitly checked that the deformed solutions satisfy the BPS equations for type IIB theory, since the direction corresponding to the $U(1)$ R-symmetry is not involved in these deformations, we expect that the deformed solutions should preserve a fraction of supersymmetry everywhere.

This paper is organized as follows. In the next section, we consider deformations of a certain region along Coulomb branch of $\mathcal{N} = 4$ super Yang Mills theory at zero temperature, described by D3-branes distributed uniformly on an ellipsoidal shell [18]. Although this Coulomb branch is maximally supersymmetric, conformal symmetry is broken— to be regained only at the UV fixed point. This static gravity solution arises in the extremal limit of rotating black D3-branes. The three rotational parameters ℓ_i of the rotating branes become distributional parameters of the ellipsoidal shell. While a stack of coincident D3-branes corresponds to the superconformal origin of the moduli space, moving some of these branes away from each other corresponds to giving expectation values to certain scalar fields. The marginal deformation of $AdS_5 \times S^5$ considered in [4] lies in the UV limit of the deformation of this solution. The introduction of scalar expectation values does not change the superpotential. After the deformation, the superpotential along the Coulomb branch flow is still the same as at the UV fixed point. That is, the deformation yields phases of chiral superfields in the superpotential.

⁴Countably infinite classes of seven-dimensional Sasaki-Einstein spaces have been found in [14, 15, 9].

In section 3, we consider deformations of twisted field theories [19]. These theories are twisted since they live on a curved space and are coupled to an $SO(6)$ gauge field. They are described by D3-branes wrapped on a (quotiented) hyperbolic 2-cycle [20]. The corresponding geometry is nonsingular and remains so after the deformations. In the UV limit, the initial geometry is locally $AdS_5 \times S^5$. The dual field theory exhibits a flow “across dimensions” from a four-dimensional superconformal field theory to a two-dimensional superconformal field theory. While the initial two-dimensional field theory has either $(4, 4)$ or $(2, 2)$ supersymmetry, depending on how the 2-cycle is embedded within the rest of the internal space, the deformed two-dimensional field theory has $(2, 2)$ supersymmetry.

In section 4, we deform various other solutions which are asymptotically locally $AdS_5 \times S^5$. All of these solutions were initially obtained in five-dimensional gauged supergravity and then lifted to ten-dimensional type IIB theory. In five dimensions, these solutions correspond to a supersymmetric AdS_5 black hole [21, 22], a deformation (in the sense of adding $U(1)$ gauge fields) of AdS_5 [23, 24], an AdS_5 magnetic string [25, 26] and a flux-brane⁵ [27]. As we will discuss, some of these solutions are related to the solutions considered in sections 2 and 3 and have interesting holographic interpretations.

In section 5, we will discuss deformations which do not preserve any supersymmetry. In this case, there are two possibilities: either the initial solution has supersymmetries which are broken by the deformations or else the initial solution is itself nonsupersymmetric. As examples of both scenarios, we consider a couple of generalizations of the $\mathcal{N} = 4$ Coulomb branch flow [18] discussed in section 2. First we discuss an $\mathcal{N} = 1$ flow [30, 31] which asymptotes to the $\mathcal{N} = 4$ Coulomb branch flow in the UV limit. This solution also encompasses the $\mathcal{N} = 1$ Pilch-Warner flow [32, 33]. Since the deformation involves the direction associated with the $U(1)$ R-symmetry, the resulting solution is nonsupersymmetric. To illustrate the second possibility, we consider a nonsupersymmetric generalization of the $\mathcal{N} = 4$ Coulomb branch flow, which corresponds to the gauge theory being at finite temperature [18]. In this case, there are actually three directions which can be used for the deformations. After pairing these directions up with a chain of dualities and shifts, this yields a solution with six deformation parameters which, in the UV limit, reduces to the solution discussed in [34, 35, 36]. There, it was found that a three-parameter case of the latter solution has an associated string sigma model which is integrable. Nonsupersymmetric deformations have also considered in [13]. Since these deformed backgrounds were obtained from solutions of type IIB supergravity by a string of dualities and shifts, we expect that they are also solutions to the type IIB equations of motion. However, we have not explicitly checked this.

⁵This solution was recently renamed an anti-bubble in [28].

In section 6, we present some suggestions for further directions. We have also included some details of the more complicated solutions in the appendices.

2 Supersymmetric deformation of the Coulomb branch flow

Rotating black D3-branes⁶ have three independent $SO(6)$ rotational parameters ℓ_i . In the extremal limit, rotating black D3-branes reduce to a static and continuous D3-brane distribution. The three parameters ℓ_i now specify the ellipsoidal shape of the D3-brane distribution [18]. In five dimensions, the near-horizon limit of this solution corresponds [38, 17] to a supersymmetric, charged singularity which is asymptotically AdS_5 ⁷ [39]. The ellipsoidal shell containing the uniform distribution of D3-branes tends to be singular, which could represent a phase transition at the associated scale of the RG flow [18]. Therefore, the supergravity description breaks down around this region.

To better understand this configuration, one can first consider the superconformal origin of the moduli space, which corresponds to a single stack of coincident D3-branes. Since these D3-branes are each in a BPS state, moving them away from each other does not require any additional energy. Therefore, any distribution of these branes is still maximally supersymmetric. However, with the exception of the special case in which the stack is expanded out to a perfectly spherically symmetric distribution, the conformal invariance of the theory is generally broken. Nevertheless, the geometry asymptotes to $AdS_5 \times S^5$ at the UV fixed point, where conformal symmetry is regained. Moving these branes away from each other has another interesting effect. That is, at the superconformal point there are six scalar fields which transform in the vector representation of the $SO(6)$ R-symmetry and the adjoint representation of $SU(N)$, where N is the number of D3-branes. As branes are moved, these scalars take on expectation values. Therefore, the resulting supergravity solution corresponds to a certain region along Coulomb branch of $\mathcal{N} = 4$ super Yang Mills theory at zero temperature [18].

The introduction of scalar expectation values does not change the superpotential of the theory. After the deformation, the superpotential along the Coulomb branch flow is the same as the marginal deformation of $AdS_5 \times S^5$ [4], which lies at the UV fixed point. Namely, the deformed superpotential is given by

$$W_{\text{marg}} = \text{Tr} (\Phi_1 \Phi_2 \Phi_3 - q \Phi_2 \Phi_1 \Phi_3), \quad q^n = 1, \quad (2.1)$$

⁶Spinning branes were first constructed in [37].

⁷This solution is usually referred to as a charged AdS_5 black hole, which is a misnomer since there is no event horizon around the singularity.

which in the $\mathcal{N} = 1$ language is a q -deformation of the superpotential preserving $\mathcal{N} = 1$ supersymmetry [5] (See also [40, 41, 42]). The moduli space of vacua depends on q and the super Yang-Mills coupling is a function of q . For $q = 1$, the $\mathcal{N} = 4$ theory can be described by an $\mathcal{N} = 1$ theory with three adjoint chiral superfields Φ_i coupled through the superpotential $W_{\text{marg}}(q = 1)$.

We will now consider the near-horizon limit of the aforementioned distribution of D3-branes in detail. The metric and 4-form potential can be written as⁸ [18]

$$\begin{aligned} ds_{10}^2 &= f^{-1/2} \frac{r^2}{R^2} (-dt^2 + dx_1^2 + dx_2^2 + dx_3^2) + f^{-1/2} \frac{R^2}{r^2} \frac{dr^2}{L_1 L_2 L_3} \\ &+ f^{1/2} R^2 \left[k^{-1} [d\alpha + \frac{k}{4} (L_2 - L_3) s_{2\alpha} s_{2\theta} d\theta]^2 + k f^{-1} s_\alpha^2 d\theta^2 \right. \\ &+ \left. L_1 c_\alpha^2 (d\psi - d\varphi_2)^2 + L_2 s_\alpha^2 s_\theta^2 (d\psi - d\varphi_1)^2 + L_3 s_\alpha^2 c_\theta^2 (d\psi + d\varphi_1 + d\varphi_2)^2 \right], \\ F_{(5)} &= dC_{(4)} + *dC_{(4)}, \quad C_{(4)} = \left(f^{-1} \frac{r^4}{R^4} - 1 \right) dt \wedge dx_1 \wedge dx_2 \wedge dx_3, \end{aligned} \quad (2.2)$$

where $R = (4\pi g_s N)^{1/4}$. We have introduced the following quantities

$$f^{-1} = (L_1^{-1} c_\alpha^2 + L_2^{-1} s_\alpha^2 s_\theta^2 + L_3^{-1} s_\alpha^2 c_\theta^2) L_1 L_2 L_3, \quad g^{-1} = L_2 s_\theta^2 + L_3 c_\theta^2, \quad k^{-1} = L_1 s_\alpha^2 + g^{-1} c_\alpha^2, \quad (2.3)$$

and

$$L_i = 1 + \frac{\ell_i^2}{r^2}. \quad (2.4)$$

This geometry approaches $AdS_5 \times S^5$ as $r \rightarrow \infty$, in which case f, g, k , and L_i go to 1. The vector representation **6** of $SO(6)$ with respect to the full Cartan subgroup $SO(2)^3$ decomposes as $(\mathbf{2}, \mathbf{1}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{2}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{1}, \mathbf{2})$ [43]. The $SO(2)^3$ isometry group of above geometry corresponds to $U(1)^2$ global symmetry generated by φ_1 and φ_2 and $U(1)_R$ R -symmetry generated by ψ of the dual field theory.

Let us write the solution in the form given by equation (A.7) in [4]. That is, the metric is written such that the T^2 symmetry along the φ_1 and φ_2 directions is explicit:

$$\begin{aligned} ds_{IIB}^2 &= F \left[\frac{1}{\sqrt{\Delta}} (D\varphi_1 - C D\varphi_2)^2 + \sqrt{\Delta} D\varphi_2^2 \right] + F^{-1/3} g_{\mu\nu} dx^\mu dx^\nu, \\ F_{(5)} &= R^{-4} d(f^{-1} r^4) dt \wedge dx_1 \wedge dx_2 \wedge dx_3 + d\mathcal{A}_{(2)} \wedge D\varphi_1 \wedge D\varphi_2, \end{aligned} \quad (2.5)$$

⁸We have used the solution given in [30] in terms of the notations in [18], except that we have taken $\theta \rightarrow \alpha + \pi/2$, $\phi \rightarrow \theta$ in order to match the deformation of $AdS_5 \times S^5$ in [4] in the UV limit. Note that the expression in [30] corrects a mistake from that of [18]. Also, the ψ rotation generates $U(1)_R$ symmetry as in [4]. We use the simple notation $c_\theta \equiv \cos \theta$ and $s_\theta \equiv \sin \theta$ and so on.

where

$$\begin{aligned}
F^{-1/3} g_{\mu\nu} dx^\mu dx^\nu &= f^{-1/2} \frac{r^2}{R^2} (-dt^2 + dx_1^2 + dx_2^2 + dx_3^2) + f^{-1/2} \frac{R^2}{r^2} \frac{dr^2}{L_1 L_2 L_3} \\
&+ f^{1/2} R^2 \left[k^{-1} [d\alpha + \frac{k}{4} (L_2 - L_3) s_{2\alpha} s_{2\theta} d\theta]^2 \right. \\
&\left. + k f^{-1} s_\alpha^2 d\theta^2 + \frac{9}{4} g h L_1 L_2 L_3 s_{2\alpha}^2 s_{2\theta}^2 d\psi^2 \right]. \tag{2.6}
\end{aligned}$$

We have explicitly taken the Hodge dual in $F_{(5)}$ to find

$$\begin{aligned}
d\mathcal{A}_{(2)} &= \frac{3}{4} R^4 \frac{f^2}{r} s_{2\alpha} s_{2\theta} d\psi \wedge \left(\frac{L_1 L_2 L_3}{r^2} \partial_r (f^{-1} r^4) s_\alpha^2 d\alpha \wedge d\theta + k f^{-1} \partial_\alpha (f^{-1}) s_\alpha^2 dr \wedge d\theta \right. \\
&\left. + \left[\frac{L_3 - L_2}{4} s_{2\alpha} s_{2\theta} \partial_\alpha (f^{-1}) + \frac{1}{k} \partial_\theta (f^{-1}) \right] dr \wedge [d\alpha + \frac{k}{4} (L_2 - L_3) s_{2\alpha} s_{2\theta} d\theta] \right). \tag{2.7}
\end{aligned}$$

Note that $d\mathcal{A}_{(2)}$ reduces to $12R^4 c_\alpha s_\alpha^3 s_\theta c_\theta d\psi \wedge d\alpha \wedge d\theta$ in asymptotic limit $r \rightarrow \infty$, which is consistent with how [4] expressed the potential $C_{(4)}$ in the $AdS_5 \times S^5$ solution. In the above, we have introduced the connection one-forms

$$\begin{aligned}
D\varphi_1 &\equiv d\varphi_1 + A_\psi^1 d\psi, & D\varphi_2 &\equiv d\varphi_2 + A_\psi^2 d\psi, \\
A_\psi^1 &= -g L_3 c_\theta^2 (2g L_2 L_3 s_\alpha^2 s_{2\theta}^2 - 4L_1 c_\alpha^2) h + g (L_3 c_\theta^2 - L_2 s_\theta^2), \\
A_\psi^2 &= (2g L_2 L_3 s_\alpha^2 s_{2\theta}^2 - 4L_1 c_\alpha^2) h, \tag{2.8}
\end{aligned}$$

where

$$h^{-1} = 4L_1 c_\alpha^2 + g L_2 L_3 s_\alpha^2 s_{2\theta}^2. \tag{2.9}$$

By comparing the metric (2.2) with (2.5), the metric functions are given by

$$F(r, \theta, \alpha) = \frac{f^{1/2} R^2 s_\alpha}{2\sqrt{gh}}, \quad \Delta(r, \theta, \alpha) = \frac{g}{4h s_\alpha^2}, \quad C(r, \theta) = -g L_3 c_\theta^2 \tag{2.10}$$

where the functions f, g , and h are given in (2.3) and (2.9).

We will now use the deformation formulae presented in the appendix of [4]. The $SL(3, R)$ transformation can be described by the matrix

$$\Lambda = \begin{pmatrix} 1 & \gamma & 0 \\ 0 & 1 & 0 \\ 0 & \sigma & 1 \end{pmatrix}. \tag{2.11}$$

This transformation is parameterized by the real parameters γ and σ , where $\beta = \gamma - \tau_s \sigma$ is related to the aforementioned q by $q = e^{2\pi i \beta}$. In general, τ_s is a complex structure parameter that is related to the gauge coupling and theta parameter of the dual gauge theory.

The transformation of various scalar fields can be obtained by constructing 3×3 matrix g^T [4] where

$$g^T = \begin{pmatrix} e^{-\phi/3} F^{-1/3} & 0 & 0 \\ 0 & e^{-\phi/3} F^{2/3} & 0 \\ 0 & 0 & e^{2\phi/3} F^{-1/3} \end{pmatrix} \begin{pmatrix} 1 & B_{12} & 0 \\ 0 & 1 & 0 \\ \chi_{\text{axion}} & -C_{12} + \chi_{\text{axion}} B_{12} & 1 \end{pmatrix}. \quad (2.12)$$

We choose the dilaton and axion fields to be vanishing: $\phi_{\text{dilaton}} = \chi_{\text{axion}} = 0$. Then τ_s is simply i and β reduces to $\gamma - i\sigma$. However, in general the dilaton and axion can be specified by nonvanishing constants.

Under the $SL(3, R)$ transformation $M = gg^T$ goes to $M' = \Lambda M \Lambda^T$ and the corresponding g^T transforms as [4]

$$g^T = \begin{pmatrix} \frac{1}{\sqrt{GH}F^{1/3}} & 0 & 0 \\ 0 & \sqrt{G}F^{2/3} & 0 \\ 0 & 0 & \sqrt{H}F^{-1/3} \end{pmatrix} \begin{pmatrix} 1 & \gamma G F^2 & 0 \\ 0 & 1 & 0 \\ \gamma\sigma F^2 H^{-1} & \sigma F^2 H^{-1} & 1 \end{pmatrix}, \quad (2.13)$$

where

$$H \equiv 1 + \sigma^2 F^2, \quad G^{-1} \equiv 1 + (\gamma^2 + \sigma^2) F^2. \quad (2.14)$$

Also, the $\mathcal{A}_{(2)}$ term in the initial five-form field strength induces additional terms in the deformed two-forms B and $C^{(2)}$ as follows [4]:

$$\begin{pmatrix} \frac{1}{2} \tilde{c}_{\mu\nu} dx^\mu \wedge dx^\nu \\ \mathcal{A}_{(2)} \\ \frac{1}{2} \tilde{b}_{\mu\nu} dx^\mu \wedge dx^\nu \end{pmatrix} \longrightarrow \Lambda \begin{pmatrix} \frac{1}{2} \tilde{c}_{\mu\nu} dx^\mu \wedge dx^\nu \\ \mathcal{A}_{(2)} \\ \frac{1}{2} \tilde{b}_{\mu\nu} dx^\mu \wedge dx^\nu \end{pmatrix}. \quad (2.15)$$

By comparing (2.12) with (2.13), one can read off all the deformed fields. By reading off the deformed fields from (A.15) of [4], we find that the resulting solution can be written (with the metric expressed in the string frame) as

$$\begin{aligned} ds_{IB}^2 &= G\sqrt{H} F \left[\frac{1}{\sqrt{\Delta}} (D\varphi_1 - C D\varphi_2)^2 + \sqrt{\Delta} D\varphi_2^2 \right] + \sqrt{H} F^{-1/3} g_{\mu\nu} dx^\mu dx^\nu, \\ B &= \gamma F^2 G D\varphi_1 \wedge D\varphi_2 - \sigma \mathcal{A}_{(2)}, \quad C^{(2)} = -\sigma F^2 G D\varphi_1 \wedge D\varphi_2 - \gamma \mathcal{A}_{(2)}, \\ e^{2\phi} &= H^2 G, \quad \chi = \gamma\sigma F^2 H^{-1}, \end{aligned} \quad (2.16)$$

and

$$F_{(5)} = dC_{(4)} + *dC_{(4)}, \quad C_{(4)} = \left(f^{-1} \frac{r^4}{R^4} - 1 \right) dt \wedge dx_1 \wedge dx_2 \wedge dx_3, \quad (2.17)$$

where the metric functions are given by (2.10), the covariant derivatives are defined in (2.8), and $d\mathcal{A}_{(2)}$ is given by (2.7). Note that the Hodge dual in $F_{(5)}$ is now taken with respect to the deformed metric.

As can be seen from the above solution, the β -deformation results in turning on the complex 3-form field strength, which is given by $d(B + iC_{(2)}) = \beta d(F^2 G D\varphi_1 \wedge D\varphi_2 - i\mathcal{A}_{(2)})$. This supports D5 and NS5-branes. As discussed in [4], in order for the 5-brane charges to be quantized, β should take on fractional values. Also, since the dilaton is no longer constant, the corresponding super Yang-Mills coupling constant runs along the flow.

One can check that the deformed geometry is regular, other than at the singular shell of D3-branes. In fact, it is easy to see that the deformations do not result in additional power-law type curvature singularities to the geometry, since there are no new poles in the metric. This is a general statement that applies to this choice of $SL(3, R)$ deformations [4], including all of the examples discussed in this paper. However, since the initial solution has poles in the metric, one has to check that the deformations do not result in conical singularities. For the present solution, the metric components either blow up or vanish when either $s_\alpha \rightarrow 0$ or $c_\alpha \rightarrow 0$, $s_{2\theta} \rightarrow 0$. In either case F vanishes, which means that both G and H go to unity. This guarantees that the deformed metric reduces to the original metric near the poles, so that the regularity conditions on coordinate periods are unchanged. We have also checked that all of the fields are single-valued at these poles. These conclusions also apply to the other examples to be discussed.

3 Supersymmetric deformations of twisted field theories

We will now consider deformations of field theories which live on D3-branes wrapped on H^2 , where H^2 is a (quotiented) hyperbolic 2-cycle [20]. These theories are coupled to an $SO(6)$ gauge field and are known as twisted field theories [19]. In some cases, the corresponding geometry in type IIB supergravity exhibits a smooth flow from an asymptotically locally $AdS_5 \times S^5$ geometry to a warped product of $AdS_3 \times H^2$ and an internal space. This corresponds to an RG flow “across dimensions” from a four-dimensional superconformal field theory to a two-dimensional superconformal field theory. The 2-cycle can be embedded within the rest of the internal space in a number of different ways, which corresponds to a certain choice of external $SO(6)$ gauge fields on the worldvolume theory. This dictates the amount of supersymmetry preserved by the two-dimensional theory in the IR. We will consider a general form of the solution as it was presented in [20] which encompasses the cases in which the two-dimensional field theory preserves either $(4, 4)$ or $(2, 2)$ supersymmetry.

First, we will turn to some of the details of the system in five-dimensional gauged supergravity, which involves the metric, a $U(1)$ gauge field and a scalar field which corresponds to

turning on an operator in the **20** of $SO(6)$. Consider an ansatz of the form

$$\begin{aligned} ds_5^2 &= e^{2f(r)}(dr^2 + dz^2 - dt^2) + \frac{e^{2g(r)}}{y^2}(dx^2 + dy^2), \\ A_{(1)}^1 &= A_{(1)}^2 = a \frac{dx}{y}, \quad A_{(1)}^3 = (1 - 2a) \frac{dx}{y}, \end{aligned} \quad (3.1)$$

along with a scalar field φ . The supersymmetry variation equations of the fermionic fields imply that f , g and φ obey a system of first-order differential equations, which are written explicitly in [20]. In particular, it was found that taking $a = 0$ or $1/2$ leads to cases in which the two-dimensional field theory has $(4, 4)$ or $(2, 2)$ supersymmetry, respectively.

The exact solution found in [20] with $(4, 4)$ supersymmetry actually flows to a singularity in the IR. Also, the equations were only partially solved for the $(2, 2)$ supersymmetric case which smoothly interpolates between local AdS_5 and $AdS_3 \times H^2$. Smooth interpolations between these conformal fixed points have been found numerically [44]. Solutions which interpolate between local AdS_5 and $AdS_3 \times S^2$ have also been found, although the dual gauge theory has not been discussed. For the case in which the three $U(1)$ gauge fields are all equally-charged and the scalar is constant, there is a exact solution [25, 26]. However, since some fields acquire fractional spins, it is not clear if this solution actually makes any sense [20]. Nevertheless, since deformations of the types of solutions discussed in this section have similar form to this exact solution, we provide details of this in section 4.4.

The system described by (3.1) can be lifted to type IIB theory using the dimensional reduction ansatz in [17]. We can express the resulting ten-dimensional metric in T^2 form as⁹

$$ds_{10}^2 = \sqrt{\tilde{\Delta}} ds_5^2 + F \left[\frac{1}{\sqrt{\tilde{\Delta}}} (D\varphi_1 - C D\varphi_2)^2 + \sqrt{\tilde{\Delta}} D\varphi_2^2 \right] \quad (3.2)$$

$$+ \frac{R^2}{\sqrt{\tilde{\Delta}}} e^{-\varphi} \left[(c_\alpha^2 + e^{3\varphi} s_\alpha^2) d\alpha^2 + s_\alpha^2 d\theta^2 + \frac{9}{16} k e^{3\varphi} s_{2\alpha}^2 s_{2\theta}^2 \left(d\psi + \frac{1}{3R} \frac{dx}{y} \right)^2 \right], \quad (3.3)$$

where the various metric functions are

$$\begin{aligned} \tilde{\Delta}(r, \alpha) &= e^{-\varphi} s_\alpha^2 + e^{2\varphi} c_\alpha^2, \quad k(r, \theta, \alpha)^{-1} = e^{3\varphi} c_\alpha^2 + s_\alpha^2 c_\theta^2 s_\theta^2, \\ \Delta(r, \theta, \alpha)^{-1} &= k s_\alpha^2, \quad F(r, \theta, \alpha) = \frac{R^2}{\sqrt{k \tilde{\Delta}}} e^{-\varphi} s_\alpha, \quad C(\theta) = -c_\theta^2, \end{aligned} \quad (3.4)$$

and the connection one-forms are

$$\begin{aligned} D\varphi_i &= d\varphi_i + A_\psi^i \left(d\psi + \frac{a_i}{R} \frac{dx}{y} \right), \\ A_\psi^1 &= c_{2\theta} - c_\theta^2 A_\psi^2, \quad A_\psi^2 = (2s_\alpha^2 s_\theta^2 c_\theta^2 - e^{3\varphi} c_\alpha^2) k, \\ a_1 &= \frac{a c_{2\theta} - a_2 c_\theta^2 A_\psi^2}{A_\psi^1}, \quad a_2 = \frac{2a - k e^{3\varphi} c_\alpha^2}{A_\psi^2}. \end{aligned} \quad (3.5)$$

⁹Note that $R = g^{-1}$, where g is the cosmological parameter written in [17].

The self-dual five-form field strength is given by

$$\begin{aligned}
F_{(5)} &= -\frac{2}{R}[e^{-2\varphi}s_\alpha^2 + e^\varphi(1+c_\alpha^2)]\frac{e^{3f+2g}}{y^2}dr \wedge dz \wedge dt \wedge dx \wedge dy \\
&+ \frac{3}{2}R\frac{e^{2g+f}}{y^2}(\partial_r\varphi)s_{2\alpha}dz \wedge dt \wedge dx \wedge dy \wedge d\alpha \\
&+ \frac{R^2}{2}e^{3f-2g}\left[a e^{2\varphi}(s_{2\alpha}c_\theta^2 d\alpha - s_\alpha^2 s_{2\theta} d\theta) \wedge \left(d\psi + d\varphi_1 + d\varphi_2 + \frac{a}{R}\frac{dx}{y}\right) \right. \\
&+ a e^{2\varphi}(s_{2\alpha}s_\theta^2 d\alpha + s_\alpha^2 s_{2\theta} d\theta) \wedge \left(d\psi - d\varphi_1 + \frac{a}{R}\frac{dx}{y}\right) \\
&+ \left.(2a-1)e^{-4\varphi}s_{2\alpha}d\alpha \wedge \left(d\psi - d\varphi_2 + \frac{1-2a}{R}\frac{dx}{y}\right)\right] \wedge dr \wedge dz \wedge dt + \text{dual}.
\end{aligned} \tag{3.6}$$

Computing the Hodge dual terms in $F_{(5)}$ explicitly gives the term that is relevant for computing contributions to the $C^{(2)}$ and B fields in the deformed solution. This term is given by

$$F_{(5)} = d\mathcal{A}_{(2)} \wedge D\varphi_1 \wedge D\varphi_2 + \dots, \tag{3.7}$$

where $d\mathcal{A}_{(2)}$ has the form

$$\begin{aligned}
d\mathcal{A}_{(2)} &= \left(F_{\alpha\theta\psi}d\alpha \wedge d\theta + F_{r\theta\psi}dr \wedge d\theta + F_{y\theta\psi}dy \wedge d\theta + a F_{y\alpha\psi}dy \wedge d\alpha\right) \wedge \left(d\psi + \frac{1}{3R}\frac{dx}{y}\right) \\
&+ \left(F_{xy\theta}d\theta + a F_{xy\alpha}d\alpha\right) \wedge dx \wedge dy.
\end{aligned} \tag{3.8}$$

The $F_{\mu\nu\lambda}$ coefficients are presented in their full glory in appendix A. In the above expression for $F_{(5)}$ and for the rest of the paper, the $\dots$ represent terms which do not include $D\varphi_1 \wedge D\varphi_2$ in the wedge product. These terms do not contribute to the final two-form fields, as can be seen from the deformation formulae in [4].

The resulting deformed solution is given by the expression (2.16) along with the 5-form in (3.6). The 5-form is the same as before the deformation except that the Hodge dual is taken with respect to the deformed metric.

4 Further examples of supersymmetric deformations

We will now consider various backgrounds in type IIB theory which are asymptotically locally (and in some cases globally) $AdS_5 \times S^5$. On general grounds, one expects that these gravity solutions are dual to four-dimensional super Yang-Mills theories which exhibit RG flows from an $\mathcal{N} = 4$ superconformal fixed point in the UV limit. The deformed theories have an $\mathcal{N} = 1$ superconformal fixed point in the UV limit. In all of these examples, the dual field theory

lives on a curved background. In sections 4.1 and 4.2, the dual field theory lives on $\mathbf{R} \times S^3$. In sections 4.3 and 4.4, the field theory is on $\text{Minkowski}_2 \times H^2$ (like the twisted theories of section 3) and $AdS_3 \times S^1$, respectively.

4.1 Supersymmetric AdS_5 black hole

As we already mentioned, the maximally-supersymmetric continuous distributions of D3-branes discussed in section 2 tend to be singular along the surface containing the branes. This means that the corresponding solution in five-dimensional gauged supergravity is actually a naked singularity, rather than a black hole. On the other hand, rotating black D3-branes do correspond to black holes in five dimensions, but these are not supersymmetric. It has only been quite recently that supersymmetric AdS_5 black holes have been known [21]. It is a requirement that there is nonvanishing angular momentum. These solutions were coupled to arbitrarily many abelian vector multiplets in [22].

For the case in which there is a single gauge field¹⁰, the supersymmetric AdS_5 black hole solution is given [21]

$$\begin{aligned} ds_5^2 &= -\mathcal{F}^2(dt + \Psi \sigma_3)^2 + \frac{R^2 dr^2}{\mathcal{F}^2(R^2 + 2\omega^2 + r^2)} + \frac{r^2}{4} \left[\sigma_1^2 + \sigma_2^2 + \left(\frac{R^2 + 2\omega^2 + r^2}{R^2} \right) \sigma_3^2 \right], \\ A_{(1)} &= \frac{1}{R} \left[\mathcal{F} dt + \frac{\eta \omega^4}{4Rr^2} \sigma_3 \right], \end{aligned} \quad (4.1)$$

where

$$\mathcal{F} = 1 - \frac{\omega^2}{r^2}, \quad \Psi = -\frac{\eta r^2}{2R} \left(1 + \frac{2\omega^2}{r^2} + \frac{3\omega^4}{2r^2(r^2 - \omega^2)} \right). \quad (4.2)$$

$\eta = \pm 1$ and ω is a constant. The σ_i are left-invariant $SU(2)$ one-forms given by

$$\sigma_1 = s_{\phi_3} d\phi_1 - s_{\phi_1} c_{\phi_3} d\phi_2, \quad \sigma_2 = c_{\phi_3} d\phi_1 + s_{\phi_1} s_{\phi_3} d\phi_2, \quad \sigma_3 = d\phi_3 + c_{\phi_1} d\phi_2. \quad (4.3)$$

It has been shown that the above geometry is globally AdS_5 at large distance [21].

This solution can be lifted to ten-dimensional type IIB theory using the ansatz in [17], which yields

$$\begin{aligned} ds_{10}^2 &= ds_5^2 + R^2 [d\alpha^2 + s_\alpha^2 d\theta^2 + s_\alpha^2 c_\theta^2 (d\psi + d\varphi_1 + d\varphi_2 + A_{(1)})^2 \\ &\quad + c_\alpha^2 (d\psi - d\varphi_2 + A_{(1)})^2 + s_\alpha^2 s_\theta^2 (d\psi - d\varphi_1 + A_{(1)})^2], \\ F_{(5)} &= \frac{r^3}{2R} dt \wedge dr \wedge \sigma_1 \wedge \sigma_2 \wedge \sigma_3 + \frac{\omega^2}{8r^3} [d(\mu_1^2) \wedge (d\psi - d\varphi_2 + A_{(1)}) \\ &\quad + d(\mu_3^2) \wedge (d\psi - d\varphi_1 + A_{(1)}) + d(\mu_2^2) \wedge (d\psi + d\varphi_1 + d\varphi_2 + A_{(1)})] \\ &\quad \wedge [r^3(R^2 + 2\omega^2 + r^2) \sigma_1 \wedge \sigma_2 \wedge \sigma_3 + R\mathcal{F}^2 r (4R\Psi + \eta\omega^2) (dt + \Psi \sigma_3) \wedge \sigma_1 \wedge \sigma_2 \\ &\quad + 2R\eta\omega^2 dt \wedge dr \wedge \sigma_3] + \text{dual}, \end{aligned} \quad (4.4)$$

¹⁰We absorbed a $1/R$ factor in the gauge field for typographical convenience.

where

$$\mu_1 = c_\alpha \quad \mu_2 = s_\alpha c_\theta, \quad \mu_3 = s_\alpha s_\theta. \quad (4.5)$$

It has been shown that this solution preserves two supersymmetries in type IIB theory [24]. At large distance, this solution is dual to an $\mathcal{N} = 4$ superconformal field theory. It is not known if this complete solution can arise in the near-horizon limit of an asymptotically flat D3-brane configuration. However, if it does, then the D3-branes would exhibit rotation in both the transverse space as well as on the worldvolume.

The metric (4.4) can be expressed as

$$\begin{aligned} ds_{10}^2 = & F \left[\frac{1}{\sqrt{\Delta}} (D\varphi_1 - CD\varphi_2)^2 + \sqrt{\Delta} D\varphi_2^2 \right] + ds_5^2 \\ & + R^2 \left[d\alpha^2 + s_\alpha^2 d\theta^2 + \left(\frac{9c_\alpha^2 s_\alpha^2 s_{2\theta}^2}{4c_\alpha^2 + s_\alpha^2 s_{2\theta}^2} \right) (d\psi + A_{(1)})^2 \right], \end{aligned} \quad (4.6)$$

where the 5-metric is given by (4.1) and the metric functions are given by

$$F(\theta, \alpha) = R^2 s_\alpha \sqrt{c_\alpha^2 + s_\alpha^2 s_{2\theta}^2 c_\theta^2}, \quad \Delta(\theta, \alpha) = \frac{c_\alpha^2 + s_\alpha^2 s_{2\theta}^2 c_\theta^2}{s_\alpha^2}, \quad C(\theta) = -c_\theta^2. \quad (4.7)$$

We have introduced the connection one-forms

$$\begin{aligned} D\varphi_1 &\equiv d\varphi_1 + A_\psi^1 (d\psi + A_{(1)}), & D\varphi_2 &\equiv d\varphi_2 + A_\psi^2 (d\psi + A_{(1)}), \\ A_\psi^1 &= c_{2\theta} - c_\theta^2 \frac{-c_\alpha^2 + 2s_\alpha^2 s_{2\theta}^2 c_\theta^2}{c_\alpha^2 + s_\alpha^2 s_{2\theta}^2 c_\theta^2}, & A_\psi^2 &= \frac{-c_\alpha^2 + 2s_\alpha^2 s_{2\theta}^2 c_\theta^2}{c_\alpha^2 + s_\alpha^2 s_{2\theta}^2 c_\theta^2}. \end{aligned} \quad (4.8)$$

Computing the Hodge dual terms in $F_{(5)}$ explicitly gives the term of the form (3.7) which is relevant for computing the deformed two-form fields, where

$$\begin{aligned} d\mathcal{A}_{(2)} &= 6R^4 s_{2\theta} s_\alpha^3 c_\alpha d\alpha \wedge d\theta \wedge (d\psi + A_{(1)}) \\ &+ \frac{\omega^2 R^2}{48r^3} (4c_\alpha^2 + s_\alpha^2 s_{2\theta}^2) \left(\frac{2s_\alpha^2}{s_{2\theta}} [(1 + c_\theta^2) A_\psi^2 + c_{2\theta} A_\psi^1] + \frac{s_\alpha}{c_\alpha} (2A_\psi^1 + A_\psi^2) \right) \\ &\times [8R (dt + \Psi \sigma_3) \wedge dr \wedge d\alpha + 2(4R\Psi + \eta\omega^2) dr \wedge \sigma_3 \wedge d\alpha + \eta\omega^2 r \sigma_1 \wedge \sigma_2 \wedge d\alpha]. \end{aligned} \quad (4.9)$$

The resulting deformed solution is given by the expression (2.16) along with the 5-form in (4.4). The 5-form is the same as before the deformation except that the Hodge dual is taken with respect to the deformed metric.

4.2 Deformation of AdS_5

A one-parameter family of type IIB solutions that is asymptotically locally $AdS_5 \times S^5$ was found in [23, 24]. This is referred to as deformation of $AdS_5 \times S^5$ by the authors, which is

a different type of deformation than the one we are presently implementing. The metric has the form (4.4) with ds_5^2 given by

$$ds_5^2 = -\left(dt + \frac{r^2}{2R}\sigma_3 + \frac{fr^2}{V}\sigma_1\right)^2 + \frac{dr^2}{V} + \frac{r^2}{4}(\sigma_1^2 + \sigma_2^2 + V\sigma_3^2), \quad (4.10)$$

where

$$V(r) = 1 + r^2/R^2, \quad A_{(1)} = \frac{fr^2}{RV(r)}\sigma_1, \quad (4.11)$$

and the σ_i and μ_i are given by (4.3) and (4.5), respectively. There is also a self-dual five-form field given by

$$\begin{aligned} F_{(5)} &= \frac{r^3}{2R} dt \wedge dr \wedge \sigma_1 \wedge \sigma_2 \wedge \sigma_3 - \frac{fR^2}{2V^2} [d(\mu_1^2) \wedge (d\psi - d\varphi_2 + A_{(1)}) \\ &+ d(\mu_3^2) \wedge (d\psi - d\varphi_1 + A_{(1)}) + d(\mu_2^2) \wedge (d\psi + d\varphi_1 + d\varphi_2 + A_{(1)})] \\ &\wedge \left[dt \wedge \left(r^2 V \sigma_2 \wedge \sigma_3 - 2r dr \wedge \sigma_1 \right) - \sigma_1 \wedge \sigma_3 \wedge \left(\frac{r^3}{R} dr + f r^4 \sigma_2 \right) \right] + \text{dual}. \end{aligned} \quad (4.12)$$

This solution preserves two supersymmetries of type IIB theory [24]. When the parameter f vanishes, the above geometry reduces to $AdS_5 \times S^5$. For nonzero f , the geometry is locally asymptotically $AdS_5 \times S^5$, implying that the UV limit of the dual gauge theory is $\mathcal{N} = 4$ super Yang-Mills theory.

This geometry has the exotic property that, when $4R^2 f^2 > 1$, closed timelike curves are present; they are absent when $4R^2 f^2 \leq 1$ [23]. These closed timelike curves appear in the boundary, as well as the bulk, which implies that they are present on the background of the dual gauge theory. It has been shown that the holographic energy-momentum tensor remains finite, which might be interpreted as a hint that the AdS/CFT correspondence is still valid in the presence of closed timelike curves. Of course, it is certainly possible that these closed timelike curves are simply a sign that the solution is not physical past an upper bound on f [24].

The 10-dimensional metric can be written in the form (4.6) together with (4.10) and (4.11). Computing the Hodge dual terms in $F_{(5)}$ explicitly gives the relevant term (3.7) with

$$\begin{aligned} d\mathcal{A}_{(2)} &= 6R^4 s_{2\theta} s_\alpha^3 c_\alpha d\alpha \wedge d\theta \wedge (d\psi + A_{(1)}) \\ &- \frac{fR^3 r}{6V^2} (4c_\alpha^2 + s_\alpha^2 s_{2\theta}^2) (2dr \wedge \sigma_1 - rV \sigma_2 \wedge \sigma_3) \\ &\wedge \left[\frac{s_\alpha^3}{s_{2\theta}} [(1 + c_\theta^2) A_\psi^2 + c_{2\theta} A_\psi^1] d\theta + \frac{s_\alpha^2}{2c_\alpha} (2A_\psi^1 + A_\psi^2) d\alpha \right]. \end{aligned} \quad (4.13)$$

The resulting deformed solution is given by the expression (2.16) along with the 5-form in (4.12).

4.3 AdS_5 magnetic string

We will now discuss the deformations of a five-dimensional magnetic string solution which can be embedded in $\mathcal{N} = 2$ gauged supergravity¹¹ [25, 26]. This solution arises from the system studied in section 3 when the three $U(1)$ gauge fields are all equally-charged and the scalars are constant. However, as previously mentioned, in this case some fields acquire fractional spins, which might be a sign that this solution is not well-behaved [20]. Nevertheless, it is a simple and exact solution which shares some of the properties of other solutions which arise from the system discussed in section 3. The five-dimensional solution is given by

$$\begin{aligned} ds_5^2 &= \frac{r^2}{R^2} H^{3/2} (-dt^2 + dx^2) + \frac{R^2}{H^2 r^2} dr^2 + r^2 (d\phi_1^2 + \sinh^2 \phi_1 d\phi_2^2), \\ F_{(2)} &= -\frac{R}{3} \sinh \phi_1 d\phi_1 \wedge d\phi_2, \end{aligned} \quad (4.14)$$

where

$$H = 1 - \frac{R^2}{3r^2}. \quad (4.15)$$

The geometry interpolates between $AdS_3 \times H^2$, where H^2 is hyperbolic 2-plane, at short distance and locally AdS_5 at large distance. At short distance, the supersymmetry is enhanced by a factor of two [26]. The charge of the $U(1)$ gauge field is uniquely determined by a supersymmetry condition.

This solution can be lifted to type IIB theory, where it preserves two supersymmetries. Using the dimensional reduction ansatz given in [17], one can see that the ten-dimensional metric has the form (4.4), where ds_5^2 is given in (4.14) and $A_{(1)} = \frac{1}{3} \cosh \phi_1 d\phi_2$. The self-dual five-form is given by

$$\begin{aligned} F_{(5)} &= \frac{4r^3 \sqrt{H}}{R^2} \sinh \phi_1 dt \wedge dx \wedge dr \wedge d\phi_1 \wedge d\phi_2 \\ &+ \frac{\sqrt{H} R^2}{6r} dt \wedge dx \wedge dr \wedge [d(\mu_1^2) \wedge (d\psi - d\varphi_2 + A_{(1)}) \\ &+ d(\mu_3^2) \wedge (d\psi - d\varphi_1 + A_{(1)}) + d(\mu_2^2) \wedge (d\psi + d\varphi_1 + d\varphi_2 + A_{(1)})] + \text{dual}. \end{aligned} \quad (4.16)$$

where we take the Hodge dual with respect to the five-dimensional metric (4.14). Then the ten-dimensional metric can be written in the form (4.6).

Computing the Hodge dual terms in $F_{(5)}$ explicitly gives (3.7) where

$$\begin{aligned} d\mathcal{A}_{(2)} &= 6R^4 s_{2\theta} s_\alpha^3 c_\alpha d\alpha \wedge d\theta \wedge (d\psi + A_{(1)}) - \frac{R^4}{6} (4c_\alpha^2 + s_\alpha^2 s_{2\theta}^2) \\ &\times \left[\frac{s_\alpha}{c_\alpha} (2A_\psi^1 + A_\psi^2) d\alpha - \frac{s_\alpha^2}{s_{2\theta}} [c_{2\theta} A_\psi^2 + (1 + c_\theta^2) A_\psi^2] d\theta \right] \sinh \phi_1 d\phi_1 \wedge d\phi_2. \end{aligned} \quad (4.17)$$

¹¹It has been shown that this solution solves first-order equations which come from a superpotential [44].

The resulting deformed solution is given by the expression (2.16) along with the 5-form in (4.16). Like the solutions discussed in section 3, this may be dual to an four-dimensional $\mathcal{N} = 1$ superconformal field theory in the UV limit which flows to a two-dimensional (2, 2) superconformal field theory in the IR region.

4.4 AdS_5 flux-brane/anti-bubble

The magnetic string geometry of the previous subsection and some of the five-dimensional geometries discussed in section 3 are locally AdS_5 in the UV limit and $AdS_3 \times H^2$ in the IR region. In fact, the same $AdS_3 \times H^2$ fixed point can interpolate to two different asymptotically locally AdS_5 spacetimes. For the previously-discussed case, the local AdS_5 geometry has a boundary $Minkowski_2 \times H^2$. This is because there is an H^2 component of the metric, with a warping factor, throughout the flow. It is also possible to maintain an AdS_3 component, again with a warping factor, all along the flow. In this case, the warp factor depends on a “radial” coordinate which lies within the H^2 rather than AdS_3 , and the asymptotically locally AdS_5 has the boundary $AdS_3 \times S^1$. This five-dimensional solution [27] is given by

$$ds_5^2 = H dx^2 + H^{-1} dr^2 + r^2 ds_{AdS_3}^2, \quad *_5 F_{(2)} = \frac{8}{9\sqrt{3}} R^2 \epsilon_{(3)}, \quad (4.18)$$

where

$$H = \frac{r^2}{R^2} - \left(1 - \frac{4R^2}{27r^2}\right)^2 \quad (4.19)$$

and $\epsilon_{(3)}$ is the volume-form of the unit AdS_3 . This solution is related to the supersymmetric AdS_5 naked singularity discussed in section 2 via a double Wick rotation. This solution smoothly flows between the asymptotically locally AdS_5 region and $AdS_3 \times H^2$ at $r = \frac{2}{3}R$. This was initially called a flux-brane because the supporting magnetic two-form field strength $F_{(2)}$ is proportional to the entire transverse space spanned by the coordinates r and x [27]. More recently, it has been referred to as an anti-bubble [28]. It is interesting to note that, once the cosmological constant is fixed, there are no free parameters associated with the $F_{(2)}$ field strength or the AdS_3 curvature. Equivalently, the AdS_3 curvature fixes the five-dimensional cosmological constant as well as the charge associated with $F_{(2)}$.

Lifting this solution to type IIB theory with the ansatz given in [17] yields a ten-dimensional metric of the form (4.4) with ds_5^2 given by (4.18) and $A_{(1)} = \frac{R}{2r^2} dx$. The self-dual five-form is given by

$$\begin{aligned} F_{(5)} = & \frac{4r^3}{R} \epsilon_{(3)} \wedge dx \wedge dr - \frac{4}{27} R^4 \epsilon_{(3)} \wedge \left[d(\mu_1^2) \wedge (d\psi - d\varphi_2 + A_{(1)}) \right. \\ & \left. + d(\mu_3^2) \wedge (d\psi - d\varphi_1 + A_{(1)}) + d(\mu_2^2) \wedge (d\psi + d\varphi_1 + d\varphi_2 + A_{(1)}) \right] + \text{dual}, \end{aligned} \quad (4.20)$$

where the μ_i are given by (4.5). Then the 10-dimensional metric can be written in the form (4.6) with $D\varphi_i$ and A_ψ^i given by (4.8). This solution might be related to some of those recently discussed in [29].

The gravity background asymptotically has $SO(2, 4) \times SO(6)$ symmetry, where $SO(2, 4)$ describes the four-dimensional conformal symmetry of the UV fixed point with $SO(6)$ R-symmetry. Away from the UV limit, the symmetry of the gravity background is broken to $SO(2, 2) \times U(1)^4$. On the gauge theory side, this corresponds to the four-dimensional conformal symmetry breaking to $SO(2, 2)$, which is the $SL(2, R) \times SL(2, R)$ part of the three-dimensional conformal symmetry group. Also, a $U(1)^3$ of the R-symmetry group remains unbroken. There is an additional $U(1)$ corresponding to a circular direction in the background on which the field theory lives. As the IR region is approached, the four-dimensional description breaks down and is replaced by a three-dimensional description. Therefore, this gravity solution probably corresponds to a vacuum of $\mathcal{N} = 4$ Yang-Mills theory on $AdS_3 \times S^1$, for which a Kaluza-Klein mode on S^1 is excited¹². This KK mode corresponds to the gauge field in (4.18) from the point of view of five-dimensional gauged supergravity. From the ten-dimensional type IIB point of view, this corresponds to gravitational fluctuations associated with the $U(1)^3$ directions of the five-sphere. The scale of the KK mode is given by the parameter R . Also, this classical gravity description is valid so long as R is large compared to the string length.

The term in $F_{(5)}$ which is relevant for the deformation is (3.7) where

$$\begin{aligned} d\mathcal{A}_{(2)} &= 6R^4 s_{2\theta} s_\alpha^3 c_\alpha d\alpha \wedge d\theta \wedge (d\psi + A_{(1)}) \\ &- \frac{2R^5}{81r^3} (4c_\alpha^2 + s_\alpha^2 s_{2\theta}^2) \left[\frac{2s_\alpha^2}{s_{2\theta}} [c_{2\theta} A_\psi^1 + (1 + c_\theta^2) A_\psi^2] d\theta - \frac{s_\alpha}{c_\alpha} (2A_\psi^1 + A_\psi^2) d\alpha \right] \wedge dx \wedge dr. \end{aligned} \quad (4.21)$$

The resulting deformed solution is given by the expression (2.16) along with the 5-form in (4.20). The deformed dual field theory is a four-dimensional $\mathcal{N} = 1$ superconformal field theory on $AdS_3 \times S^1$ in the UV limit and flows to a two-dimensional $(2, 2)$ superconformal field theory in the IR region.

5 Nonsupersymmetric deformations

We now turn to a couple of examples of deformations which do not preserve any supersymmetry. Both of the initial solutions are generalizations of the $\mathcal{N} = 4$ Coulomb branch flow discussed in section 2. In section 5.1, we discuss an $\mathcal{N} = 1$ supersymmetric generalization

¹²We appreciate Juan Maldacena's input on this matter.

given in [30, 31]. Although the initial solution is supersymmetric, the resulting solution is nonsupersymmetric since the deformation involves the direction associated with the $U(1)$ R-symmetry. In section 5.2, we consider a case in which the initial solution itself is non-supersymmetric. In particular, we consider the near-horizon limit of a nonextremal rotating D3-brane, which corresponds to the gauge theory being at finite temperature [18].

5.1 $\mathcal{N} = 1$ Pilch-Warner flow

There is a three-parameter family of type IIB supergravity solutions which correspond to $\mathcal{N} = 1$ supersymmetric holographic RG flows [30, 31]. Two of the parameters represent independent scalar masses or vevs and the third represents a single fermion mass. The ten-dimensional metric of the full solution is given by [30, 31]

$$ds_{IIB}^2 = \Omega^2 (e^{2A(r)} dx_\mu^2 + dr^2) + \frac{L^2}{\Omega^2} ds_5^2 \quad (5.1)$$

where the warp factor is

$$\Omega(r, \theta, \phi) \equiv \sqrt{\cosh \chi} \left[\frac{1}{\rho^2} \left(\nu^2 c_\phi^2 + \frac{1}{\nu^2} s_\phi^2 \right) c_\theta^2 + \rho^4 s_\theta^2 \right]^{1/4}, \quad (5.2)$$

and the metric of the five-dimensional internal space is given by

$$\begin{aligned} ds_5^2 &= \frac{1}{\rho^4} [c_\theta^2 + \rho^6 s_\theta^2 (\nu^{-2} c_\phi^2 + \nu^2 s_\phi^2)] d\theta^2 + \rho^2 c_\theta^2 [\nu^2 c_\phi^2 + \nu^{-2} s_\phi^2] d\phi^2 \\ &- 2\rho^2 [\nu^2 - \nu^{-2}] s_\theta c_\theta s_\phi c_\phi d\theta d\phi + \rho^2 c_\theta^2 [\nu^{-2} c_\phi^2 d\varphi_1^2 + \nu^2 s_\phi^2 d\varphi_2^2] + \frac{1}{\rho^4} s_\theta^2 (d\varphi_1 + d\varphi_2 + d\varphi_3)^2 \\ &+ \frac{1}{\Omega^4} \sinh^2 \chi \cosh^2 \chi [c_\theta^2 (s_\phi^2 d\varphi_2 - c_\phi^2 d\varphi_1) + s_\theta^2 (d\varphi_1 + d\varphi_2 + d\varphi_3)]^2. \end{aligned} \quad (5.3)$$

From the five-dimensional gauged supergravity perspective, the three scalar fields $\chi(r), \nu(r)$ and $\rho(r)$ satisfy first-order equations which arise from a superpotential. The fermion mass vanishes asymptotically when $\chi = 0$, where the solution reduces to that of the near-horizon region of an ellipsoidal distribution of extremal D3-branes (2.2). This limit, which we have already discussed in section 2, is associated with the Coulomb branch flows of the $\mathcal{N} = 4$ super Yang-Mills theory. When $\nu = 1$, this system reduces to the Pilch-Warner flows [32, 33], which correspond to the Leigh-Strassler renormalization flow studied in [5]. We will focus on

this limit, in which case the solution can be written as¹³

$$\begin{aligned}
ds_{IIB}^2 &= \frac{X_1^{1/2} \cosh \chi}{\rho} (e^{2A(r)} dx_\mu^2 + dr^2) \\
&+ \frac{X_1^{1/2} \rho^3}{g^2 \cosh \chi} \left[\frac{4}{\rho^6} d\theta^2 + \frac{c_\theta^2}{X_1} (d\alpha^2 + s_\alpha^2 d\eta^2) + \frac{c_\theta^2 \cosh \chi}{X_2} (d\beta + c_\alpha d\eta)^2 \right. \\
&+ \left. \frac{s_\theta^2 \cosh \chi}{X_1^2 X_2 \rho^6} \left(2X_2 (d\phi - d\beta) + \rho^6 c_\theta^2 \sinh \chi \tanh \chi (d\beta + c_\alpha d\eta) \right)^2 \right], \\
F_{(5)} &= d^4 x \wedge (w_r dr + w_\theta d\theta) + * [d^4 x \wedge (w_r dr + w_\theta d\theta)], \\
B_{(2)}^{\text{NS}} + i C_{(2)}^{\text{RR}} &= e^{-i\phi} \left[i a_1 d\theta - a_2 (d\beta + c_\alpha d\eta) - a_3 (d\phi - d\beta) \right] \wedge (d\alpha + i s_\alpha d\eta). \quad (5.4)
\end{aligned}$$

The functions X_1 , X_2 , w_r , w_θ and a_i depend on r and θ and are given by (B.1), (B.2) and (B.3). They are written in terms of the scalar fields $\chi(r)$ and $\rho(r)$, which satisfy the supersymmetric flow equations [45] of five-dimensional gauged supergravity. These flow equations also involve the metric function $A(r)$, although no explicit closed analytic solutions are known. The dilaton-axion scalar field remains constant along this flow. The complex two-form field corresponds to non-zero fermion masses in the super Yang-Mills theory. The geometry has $SU(2) \times U(1)_\beta$ isometry, which contains a $U(1)_\eta \times U(1)_\beta$ subgroup. The $U(1)_\eta \times U(1)_\beta$ correspond to the η and β directions along which we will be applying the $SL(2, R)$ transformations.

At large distance, the solution has the maximally supersymmetric geometry $AdS_5 \times S^5$, which is dual to $\mathcal{N} = 4$ superconformal Yang-Mills theory. At short distance, the geometry is a warped product of AdS_5 and the internal space, which corresponds to the IR $\mathcal{N} = 1$ superconformal fixed point of the Leigh-Strassler flow [5].

We will now express this solution in the form given by equation (A.7) in [4], in order to apply their deformation procedure:

$$\begin{aligned}
ds_{IIB}^2 &= F \left[\frac{1}{\sqrt{\Delta}} (D\beta - C D\eta)^2 + \sqrt{\Delta} D\eta^2 \right] + \frac{e^{2\phi/3}}{F^{1/3}} \left(\frac{X_1 \cosh \chi}{g} \right)^{2/3} (c_1 c_4)^{1/6} \\
&\times \left[e^{2A(r)} dx_\mu^2 + dr^2 + \frac{\rho^4}{g^2 \cosh^2 \chi} \left(\frac{4}{\rho^6} d\theta^2 + \frac{c_\theta^2}{X_1} d\alpha^2 + c_6 d\phi^2 \right) \right], \quad (5.5) \\
B_{(2)}^{\text{NS}} &= B_{12} D\beta \wedge D\eta + (B_{1\mu} D\beta + B_{2\mu} D\eta) \wedge dx^\mu - \frac{1}{2} A_\mu^m B_{m\nu} dx^\mu \wedge dx^\nu + \frac{1}{2} \tilde{b}_{\mu\nu} dx^\mu \wedge dx^\nu, \\
C_{(2)}^{\text{RR}} &= C_{12} D\beta \wedge D\eta + (C_{1\mu} D\beta + C_{2\mu} D\eta) \wedge dx^\mu - \frac{1}{2} A_\mu^m C_{m\nu} dx^\mu \wedge dx^\nu + \frac{1}{2} \tilde{c}_{\mu\nu} dx^\mu \wedge dx^\nu.
\end{aligned}$$

¹³This form of the solution can be obtained by taking $\phi \rightarrow \phi - \beta$ in the solution of [33] in order to absorb the β dependence on two-form potential. The $U(1)_\beta$ symmetry under which Φ_1, Φ_2 and Φ_3 have charges $(1/2, 1/2, 0)$ exists if we allow the mass m of the chiral superfield Φ_3 to rotate by a phase [31]: $\Phi_1 \rightarrow e^{\frac{i}{2}\beta} \Phi_1, \Phi_2 \rightarrow e^{\frac{i}{2}\beta} \Phi_2, \Phi_3 \rightarrow \Phi_3, m \rightarrow e^{i\beta} m$. However, since the original ϕ rotation generates $U(1)_R$ symmetry, after this shift, still β rotation is related to this $U(1)_R$ symmetry. In other words, contrary to the previous case, one of the $U(1)$'s we take should contain $U(1)_R$ implying that the supersymmetry is broken completely.

The index m stands for 1 and 2 that are the coordinates β and η for two torus and the indices μ, ν stand for the remaining 8-dimensional coordinates and

$$\begin{aligned} D\beta &\equiv d\beta + A_\phi^1 d\phi, & D\eta &\equiv d\eta + A_\phi^2 d\phi, \\ A_\phi^1 &= (c_3 - c_2 c_5), & A_\phi^2 &= c_5. \end{aligned} \quad (5.6)$$

Also, the metric functions are given by

$$F(r, \theta) = \frac{(X_1 c_1 c_4)^{1/2} \rho^3}{g^2 \cosh \chi}, \quad \Delta(r, \theta) = \frac{c_4}{c_1}, \quad C(r, \theta) = -c_2. \quad (5.7)$$

The coefficient functions in $B_{(2)}^{\text{NS}}$ and $C_{(2)}^{\text{RR}}$ are given by (B.4), and the functions c_i are given in (B.5). The resulting nonsupersymmetric type IIB deformed metric in the string frame is given by

$$\begin{aligned} ds_{IIB}^2 &= G\sqrt{H} F \left[\frac{1}{\sqrt{\Delta}} (D\beta' - C D\eta')^2 + \sqrt{\Delta} D\eta'^2 \right] \\ &+ \sqrt{H} \frac{X_1^{1/2} \cosh \chi}{\rho} \left[e^{2A(r)} dx_\mu^2 + dr^2 + \frac{\rho^4}{g^2 \cosh^2 \chi} \left(\frac{4}{\rho^6} d\theta^2 + \frac{c_\theta^2}{X_1} d\alpha^2 + c_6 d\phi^2 \right) \right], \\ D\beta' &\equiv D\beta + (\gamma B_{2\mu} - \sigma C_{2\mu}) dx^\mu, & D\eta' &\equiv D\eta + (\sigma C_{1\mu} - \gamma B_{1\mu}) dx^\mu, \end{aligned} \quad (5.8)$$

and the dilaton-axion scalar fields ¹⁴ are

$$e^{2\phi} = H^2 G, \quad \chi = [\gamma \sigma (F^2 + B_{12}^2 + C_{12}^2) + \sigma B_{12} - \gamma C_{12}] H^{-1}, \quad (5.9)$$

One can also straightforwardly compute the deformed form-fields by inputting the above initial solution into the deformation formulae in [4] but the results are too long to present. At the UV fixed point, this deformed solution reduces to a particular nonsupersymmetric marginal deformation of $AdS_5 \times S^5$ which was found in [34].

5.2 Nonextremal rotating D3-brane

We now consider the nonextremal rotating D3-brane solution of type IIB theory. This solution is nonsupersymmetric and is conjectured to correspond to the Coulomb branch of a field theory at finite temperature. For the particular case in which all three angular momenta are equal,

¹⁴For nonzero B_{12} and C_{12} , the $SL(3, R)$ transformation yields $H \equiv \sigma^2 (B_{12}^2 + F^2) + (1 - \sigma C_{12})^2$, $G^{-1} \equiv \sigma^2 F^2 + \gamma^2 (F^2 + B_{12}^2) - 2\gamma B_{12}(\sigma C_{12} - 1) + (1 - \sigma C_{12})^2$. See also [4, 13, 46].

the gravity solution is given by [18]

$$\begin{aligned}
ds_{10}^2 &= \frac{Lr^2}{R^2} \left[-\frac{R^4}{L^2 r^4} dt^2 + dx_i^2 + \frac{2m}{r^4} \left(\ell^{-1} \cosh \delta dt - B_{(1)} \right)^2 \right] + \frac{R^2 L}{r^2} \frac{dr^2}{L^3 - 2m/r^4} \\
&+ R^2 [d\alpha^2 + s_\alpha^2 d\theta^2 + c_\alpha^2 (d\psi - d\varphi_2)^2 + s_\alpha^2 s_\theta^2 (d\psi - d\varphi_1)^2 + s_\alpha^2 c_\theta^2 (d\psi + d\varphi_1 + d\varphi_2)^2], \\
F_{(5)} &= dC_{(4)} + *dC_{(4)}, \\
C_{(4)} &= \left(1 - \frac{L^2 r^4}{R^4} \right) \sinh^{-1} \delta d^3 x \wedge (\cosh \delta dt - \ell B_{(1)}),
\end{aligned} \tag{5.10}$$

where

$$B_{(1)} = s_\alpha^2 c_{2\theta} d\varphi_1 + (s_\alpha^2 c_\theta^2 - c_\alpha^2) d\varphi_2 + d\psi, \tag{5.11}$$

and

$$L = 1 + \frac{\ell^2}{r^2}. \tag{5.12}$$

We have defined $R^4 \equiv 2m \sinh^2 \delta$. The quantities R , ℓ and m parameterize the D3-brane charge, angular momenta and energy above extremality, respectively. The extremal limit $m = 0$ corresponds to a static distribution of D3-branes, as was discussed in section 2. In particular, here the D3-branes would be uniformly distributed on the surface of a 5-sphere.

For $\ell^4 < \frac{8}{27}m$, there is an event horizon around the singularity. This has been interpreted as the high-density and high-temperature deconfined phase of the dual field theory. For $\ell^4 > \frac{8}{27}m$, the Hawking temperature vanishes and there is no longer an event horizon hiding the singularity. There are indications that this marks the transition from the deconfined high-density phase of the dual field theory to the Coulomb phase at finite density [47].

The metric can be written in the form given by (2.5) with

$$\begin{aligned}
F^{-1/3} g_{\mu\nu} dx^\mu dx^\nu &= c_9 dt^2 + c_7^{-1} (d\psi + c_8 dt)^2 + \frac{Lr^2}{R^2} dx_i^2 \\
&+ \frac{R^2 L}{r^2} \frac{dr^2}{L^3 - 2m/r^4} + R^2 [d\alpha^2 + s_\alpha^2 d\theta^2],
\end{aligned} \tag{5.13}$$

where

$$\begin{aligned}
F &= \frac{1}{\sqrt{c_1 c_4}}, \quad \Delta = \frac{c_1}{c_4}, \\
C &= -c_1 R^2 s_\alpha^2 \left(c_\theta^2 + \frac{L}{\sinh^2 \delta r^2} c_{2\theta} (s_\alpha^2 c_\theta^2 - c_\alpha^2) \right),
\end{aligned} \tag{5.14}$$

and

$$\begin{aligned}
D\varphi_1 &= d\varphi_1 + (c_2 + C c_5) d\psi + (c_3 + C c_6) dt, \\
D\varphi_2 &= d\varphi_2 + c_5 d\psi + c_6 dt.
\end{aligned} \tag{5.15}$$

The c_i functions are given by (B.6). The deformed solution can be expressed as (2.16) together with the $F_{(5)}$ given in (5.10). One can straightforwardly dualize $F_{(5)}$ to obtain the explicit expression for $d\mathcal{A}_{(2)}$ which is too long to present.

6 Further directions

We have considered supersymmetric deformations of various type IIB supergravity backgrounds which exhibit flows away from an asymptotically locally $AdS_5 \times S^5$ fixed point. This includes the near-horizon limit of a static ellipsoidal distribution of D3-branes which, after the deformations, corresponds to a region along the Coulomb branch of $\mathcal{N} = 1$ super Yang Mills theory. The marginal deformation of $AdS_5 \times S^5$ considered in [4] lies in the UV limit of the deformation of this solution, and the deformed superpotential does not change as one moves onto the Coulomb branch. However, there still remains a short-distance naked singularity on the surface of D3-branes in the gravity solution. This singularity can be hidden by an event horizon if the D3-branes rotate in the transverse space. This breaks supersymmetry and corresponds to giving the dual field theory at finite temperature. In order for the solution still be supersymmetric, the D3-branes must also have a specific amount of rotation within their worldvolume as well [21]. We also discuss the deformations of these solutions.

Another example of type IIB flows away from $AdS_5 \times S^5$ is provided by D3-branes wrapped on a (quotiented) hyperbolic 2-cycle [20]. This describes a twisted field theory [19] which, upon deformation, exhibits a flow “across dimensions” from an $\mathcal{N} = 1$ four-dimensional superconformal field theory to a $(2, 2)$ two-dimensional superconformal field theory. The four-dimensional field theory lives on a $Minkowski_2 \times H^2$ background. We also consider the deformation of a gravity dual which corresponds to a four-dimensional field theory which lives on $AdS_3 \times S^1$ [27], although the dual field theory has not yet been identified. Note that, after the deformations, all of these solutions still exhibit conformal fixed-points at both ends of the flow.

For all of our examples, we could have also considered dipole deformations by involving a worldvolume direction in the $SL(3, R)$ transformation [13]. In particular, for the cases in which the four-dimensional field theory lives on $Minkowski_2 \times H^2$ or $AdS_3 \times S^1$, this could involve one direction of H^2 or the S^1 direction, respectively. Then this is a dipole deformation of the four-dimensional UV field theory and a marginal deformation of the two-dimensional IR field theory. RG flows which connect these different types of deformations are rather novel.

There are a number of other type IIB solutions which are asymptotically $AdS_5 \times S^5$ and can be deformed by the technique of [4]. Such possibilities include a subclass of the bubbling

AdS solutions found in [48] and AdS_5 black holes with fermionic hair [49]. One can also apply similar deformations to the $AdS_3 \times S^3 \times S^3 \times S^1$ background. Possible holographic duals of this gravity background have been discussed in [50]. The initial two-dimensional field theory has *large* $\mathcal{N} = 4$ superconformal symmetry, due to the $SU(2) \times SU(2)$ R-symmetry. Depending on which directions of the gravity dual are used for the deformation, the deformed field theory might have $\mathcal{N} = (2, 2)$, $(0, 4)$ or *small* $(4, 4)$ supersymmetry.

We have also presented two types of nonsupersymmetric generalizations of the deformations of Coulomb branch flows. There is an $\mathcal{N} = 1$ supersymmetric flow [30, 31] which encompasses the $\mathcal{N} = 4$ Coulomb branch flow [18] and the $\mathcal{N} = 1$ Pilch-Warner flow [33]. The deformation is nonsupersymmetric, since it involves the direction associated with the $U(1)$ R-symmetry. This is clearly not the gravity dual of an $\mathcal{N} = 4$ super Yang-Mills theory which has undergone both marginal and relevant deformations [41], since the resulting solution would need to be supersymmetric. In fact, it would be interesting to find the gravity dual for this, which may require a deformation technique that does not involve T-duality. Anyway, one can also apply this deformation procedure to the $\mathcal{N} = 2$ supersymmetric RG flow solution found in [51, 52], although the result will also be nonsupersymmetric. There are also eleven-dimensional flows which can be deformed analogously, such as the ones described in [53, 54, 55].

For these deformed solutions which do not preserve any supersymmetry, it is important to study their stability. Masses should vary smoothly with the deformation parameters. Therefore, it seems reasonable to expect the lightest modes to become tachyonic first. Also, for small deformations, tachyons should be absent from the spectrum.

There is also a nonsupersymmetric generalization of the Coulomb branch flow which corresponds to the gauge theory being at finite temperature [18]. The initial solution is already nonsupersymmetric, which means there are actually three directions which can be paired up for the deformations with a chain of dualities and shifts. This yields a solution with six deformation parameters which reduces to the solution discussed in [34, 35, 36] in the UV limit. There, it was found that a three-parameter case of the latter solution has an associated string sigma model which is integrable. We do not know if this property of integrability survives away from the UV limit. The initial solution suffers from an instability for large angular momenta. Above this threshold, the D3-branes may split apart into fragments which move radially outwards and carry away some of the spin. This could correspond to a spurious gauge field vev producing a negative mass scalar term which destabilizes the moduli space of the dual gauge theory. It would be interesting to see if these deformations simply change the threshold angular momenta for stability, or if the entire solution is rendered unstable.

Acknowledgments

We would like to thank Philip Argyres, Oleg Lunin, Juan Maldacena and Carlos Nuñez for useful discussions. In particular, we appreciate the detailed explanations provided by Oleg Lunin with regards to his recent paper. The work of C.A. is supported by a grant from the Monell Foundation through the Institute for Advanced Study, and by grant No. R01-2006-000-10965-0 from the Basic Research Program of the Korea Science & Engineering Foundation. The work of J.F.V.P. is supported by DOE grant DOE-FG02-84ER-40153.

Appendix A Components of $d\mathcal{A}_{(2)}$ in section 3

The terms in $F_{(5)}$ which yield contributions to the two-forms of the deformed solution can be written in terms of $d\mathcal{A}_{(2)}$, whose components are given by

$$\begin{aligned}
F_{\alpha\theta\psi} &= \frac{6R^4}{\tilde{\Delta}^{5/2}} s_{2\theta} s_\alpha^3 c_\alpha (1 + \tilde{\Delta} e^{-2\varphi}) (c_\alpha^2 + e^{3\varphi} s_\alpha^2)^{1/2}, \\
F_{r\theta\psi} &= \frac{9}{2} R^3 s_{2\theta} s_\alpha^3 c_\alpha^2 \frac{e^{\varphi/2}}{\tilde{\Delta}^{5/4}} (c_\alpha^2 + e^{3\varphi} s_\alpha^2)^{-1/2} \varphi', \\
F_{xy\alpha} &= \frac{R^3}{3y^2} \frac{s_\alpha}{c_\alpha} \frac{2A_\psi^1 + A_\psi^2}{k\tilde{\Delta}} (c_\alpha^2 + e^{3\varphi} s_\alpha^2)^{1/2}, \\
F_{xy\theta} &= \frac{2R^3}{3y^2} \frac{s_\alpha^2}{s_{2\theta}} \frac{e^{-2\varphi}}{k\tilde{\Delta}} (c_\alpha^2 + e^{3\varphi} s_\alpha^2)^{-1/2} [a(1 - c_{2\theta} A_\psi^1 - c_\theta^2 A_\psi^2) e^{2\varphi} + (2a - 1)(1 - A_\psi^2) e^{-4\varphi}], \\
F_{y\theta\psi} &= \frac{R^4}{2y} \tilde{\Delta}^{-3/2} e^{-2g} (c_\alpha^2 + e^{3\varphi} s_\alpha^2)^{-1/2} s_\alpha^4 c_\alpha^2 s_{2\theta} [(2a + (a - 3a_1) A_\psi^1 c_{2\theta} + (a - 3a_2) A_\psi^2 c_\theta^2) a e^{2\varphi} \\
&\quad + (2(1 - 3a) + (3a_2 - a) A_\psi^2) (2a - 1) e^{-4\varphi}], \\
F_{y\alpha\psi} &= -\frac{R^4}{4} \frac{e^{-2g}}{y} s_\alpha^3 c_\alpha s_{2\theta}^2 \tilde{\Delta}^{-3/2} (c_\alpha^2 + e^{3\varphi} s_\alpha^2)^{1/2} e^{2\varphi} [2(3a_1 - a) A_\psi^1 + (3a_2 - a) A_\psi^2]. \tag{A.1}
\end{aligned}$$

Appendix B Details of the solutions in section 5

Appendix B.1 Details of the Pilch-Warner flow

The various functions in the Pilch-Warner flow of type IIB supergravity can be expressed in terms of the scalar fields $\chi(r)$ and $\rho(r)$ as [45]

$$\begin{aligned}
X_1(r, \theta) &\equiv c_\theta^2 + \rho^6 s_\theta^2, \\
X_2(r, \theta) &\equiv c_\theta^2 \operatorname{sech} \chi + \rho^6 s_\theta^2 \cosh \chi, \tag{B.1}
\end{aligned}$$

the coefficient functions for the five-form field strength are

$$\begin{aligned} w_r(r, \theta) &= \frac{g e^{4A(r)}}{8\rho^4} \cosh^2 \chi [c_\theta^2 (\cosh(2\chi) - 3) + \rho^6 (2\rho^6 s_\theta^2 \sinh^2 \chi + c_{2\theta} - 3)], \\ w_\theta(r, \theta) &= \frac{e^{4A(r)}}{8\rho^2} s_{2\theta} [2 \cosh^2 \chi + \rho^6 (\cosh(2\chi) - 3)], \end{aligned} \quad (\text{B.2})$$

and the complex two-form potential involves the coefficient functions

$$\begin{aligned} a_1(r, \theta) &= \frac{2}{g^2} c_\theta \tanh \chi, & a_2(r, \theta) &= \frac{\rho^6}{g^2 X_1} c_\theta^2 s_\theta \tanh \chi, \\ a_3(r, \theta) &= -\frac{2}{g^2 X_1} c_\theta^2 s_\theta \tanh \chi. \end{aligned} \quad (\text{B.3})$$

The complex two-form potential can be written in terms of RR and NS-NS two-form potentials $C_{(2)}^{\text{RR}}$ and $B_{(2)}^{\text{NS}}$, whose components are given by

$$\begin{aligned} (B_{12}, C_{12}) &= (a_3 - a_2) s_\alpha (s_\phi, c_\phi), & (B_{1\alpha}, C_{1\alpha}) &= (a_3 - a_2) (c_\phi, -s_\phi), \\ (B_{1\phi}, C_{1\phi}) &= (a_2 - a_3) c_5 s_\alpha (s_\phi, c_\phi), & (B_{2\phi}, C_{2\phi}) &= [a_3 + (a_3 - a_2)(c_3 - c_2 c_5)] s_\alpha (s_\phi, c_\phi), \\ (B_{2\alpha}, C_{2\alpha}) &= a_2 (-c_\phi, s_\phi) c_\alpha, & (B_{2\theta}, C_{2\theta}) &= a_1 s_\alpha (c_\phi, -s_\phi), \\ (\tilde{b}_{\phi\alpha}, \tilde{c}_{\phi\alpha}) &= -(\tilde{b}_{\alpha\phi}, \tilde{c}_{\alpha\phi}) = \frac{1}{2} [-2a_3 + (a_3 - a_2)(c_3 - c_2 c_5) - a_2 c_5 c_\alpha] (c_\phi, -s_\phi), \\ (\tilde{b}_{\theta\alpha}, \tilde{c}_{\theta\alpha}) &= -(\tilde{b}_{\alpha\theta}, \tilde{c}_{\alpha\theta}) = a_1 (s_\phi, c_\phi), & (\tilde{b}_{\theta\phi}, \tilde{c}_{\theta\phi}) &= -(\tilde{b}_{\phi\theta}, \tilde{c}_{\phi\theta}) = a_1 c_5 s_\alpha (c_\phi, -s_\phi), \end{aligned} \quad (\text{B.4})$$

where all unspecified components are zero. Notice that $C_{(2)}^{\text{RR}}(\phi) = B_{(2)}^{\text{NS}}(\phi + \pi/2)$. The c_i functions are given by

$$\begin{aligned} c_1(r, \theta) &= \frac{\cosh \chi}{X_1^2 X_2 \rho^6} (X_1^2 \rho^6 c_\theta^2 + f^2 s_\theta^2), \\ c_2(r, \theta) &= \frac{c_\theta^2 c_\alpha \cosh \chi}{X_1^2 X_2 c_1} (X_1^2 + f s_\theta^2 \sinh \chi \tanh \chi), \\ c_3(r, \theta) &= \frac{2f s_\theta^2 \cosh \chi}{X_1^2 \rho^6 c_1}, \\ c_4(r, \theta) &= \frac{c_\theta^2 s_\alpha^2}{X_1} + \frac{c_\theta^2 c_\alpha^2 \cosh \chi}{X_1^2 X_2} \left(X_1^2 + \frac{\rho^6}{4} s_{2\theta}^2 \sinh^2 \chi \tanh^2 \chi \right) - c_1 c_2^2, \\ c_5(r, \theta) &= \frac{s_{2\theta}^2 c_\alpha \sinh^2 \chi}{2X_1^2 c_4} - \frac{c_1 c_2 c_3}{c_4}, \\ c_6(r, \theta) &= \frac{4X_2 s_\theta^2 \cosh \chi}{X_1^2 \rho^6} - c_1 c_3^2 - c_4 c_5^2, \\ f(r, \theta) &\equiv \rho^6 c_\theta^2 \sinh \chi \tanh \chi - 2X_2. \end{aligned} \quad (\text{B.5})$$

Appendix B.2 Metric functions for nonextremal rotating D3-brane

The deformed near-horizon metric for a nonextremal rotating D3-brane can be expressed in terms of the functions

$$\begin{aligned}
c_1^{-1} &= R^2 s_\alpha^2 \left(1 + \frac{L}{\sinh^2 \delta r^2} s_\alpha^2 c_{2\theta}^2 \right), \\
c_2 &= c_1 R^2 s_\alpha^2 c_{2\theta} \left(1 + \frac{L}{\sinh^2 \delta r^2} \right), \\
c_3 &= -\frac{c_1 R^2 \cosh \delta}{\sinh^2 \delta r^2} s_\alpha^2 c_{2\theta}, \\
c_4^{-1} &= c_1 R^4 s_\alpha^2 \left(c_\alpha^2 + s_\alpha^2 s_\theta^2 c_\theta^2 + \frac{L}{\sinh^2 \delta r^2} (s_\alpha^2 c_\theta^2 - c_\alpha^2) [s_\alpha^2 c_\theta^2 (1 - 2c_{2\theta}) - c_\alpha^2] \right), \\
c_5 &= c_4 R^2 \left(1 + \frac{L}{\sinh^2 \delta r^2} \right) (s_\alpha^2 c_\theta^2 - c_\alpha^2 + C s_\alpha^2 c_{2\theta}), \\
c_6 &= \frac{c_4 R^2 \cosh \delta}{\sinh^2 \delta r^2} (c_\alpha^2 - s_\alpha^2 c_\theta^2 + C s_\alpha^2 c_{2\theta}), \\
c_7^{-1} &= R^2 \left(1 + \frac{L}{\sinh^2 \delta r^2} \right) (2 - 3s_\alpha^2 c_\theta^2 - C s_\alpha^2 c_{2\theta}), \\
c_8 &= -\left(\frac{R^2 \cosh \delta}{\sinh^2 \delta r^2} + c_4^{-1} c_5 c_6 + c_1^{-1} c_2 c_3 \right) c_7, \\
c_9 &= \frac{R^2}{L \sinh^2 \delta r^2} - c_1^{-1} c_3^2 - c_4^{-1} c_6^2 - c_7^{-1} c_8^2.
\end{aligned} \tag{B.6}$$

References

- [1] J. M. Maldacena, “The large N limit of superconformal field theories and supergravity,” Adv. Theor. Math. Phys. **2**, 231 (1998); Int. J. Theor. Phys. **38**, 1113 (1999) [arXiv:hep-th/9711200].
- [2] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, “Gauge theory correlators from non-critical string theory,” Phys. Lett. B **428**, 105 (1998) [arXiv:hep-th/9802109].
- [3] E. Witten, “Anti-de Sitter space and holography,” Adv. Theor. Math. Phys. **2**, 253 (1998) [arXiv:hep-th/9802150].
- [4] O. Lunin and J. Maldacena, “Deforming field theories with U(1) x U(1) global symmetry and their gravity duals,” JHEP **0505**, 033 (2005) [arXiv:hep-th/0502086].
- [5] R. G. Leigh and M. J. Strassler, “Exactly marginal operators and duality in four-dimensional N=1 supersymmetric gauge theory,” Nucl. Phys. B **447**, 95 (1995) [arXiv:hep-th/9503121].

- [6] C. Ahn and J. F. Vázquez-Poritz, “Marginal deformations with $U(1)^3$ global symmetry,” JHEP **0507**, 032 (2005) [arXiv:hep-th/0505168].
- [7] J. P. Gauntlett, D. Martelli, J. Sparks and D. Waldram, “Supersymmetric AdS(5) solutions of M-theory,” Class. Quant. Grav. **21**, 4335 (2004) [arXiv:hep-th/0402153].
- [8] J. P. Gauntlett, D. Martelli, J. Sparks and D. Waldram, “Sasaki-Einstein metrics on $S(2) \times S(3)$,” Adv. Theor. Math. Phys. **8**, 711 (2004) [arXiv:hep-th/0403002].
- [9] M. Cvetič, H. Lü, D. N. Page and C. N. Pope, “New Einstein-Sasaki Spaces in Five and Higher Dimensions,” [arXiv:hep-th/0504225].
- [10] D. Martelli and J. Sparks, “Toric Sasaki-Einstein metrics on $S(2) \times S(3)$,” Phys. Lett. **B621**, 208 (2005) [arXiv:hep-th/0505027].
- [11] I.R. Klebanov and M.J. Strassler, “Supergravity and a confining gauge theory: Duality cascades and χ SB-resolution of naked singularities,” JHEP **0008**, 052 (2000) [arXiv:hep-th/0007191].
- [12] J.M. Maldacena and C. Nuñez, “Towards the large N limit of pure $\mathcal{N} = 1$ super Yang Mills,” Phys. Rev. Lett. **86**, 588 (2001) [arXiv:hep-th/0008001].
- [13] U. Gursoy and C. Nunez, “Dipole deformations of $N = 1$ SYM and supergravity backgrounds with $U(1) \times U(1)$ global symmetry,” [arXiv:hep-th/0505100].
- [14] J. P. Gauntlett, D. Martelli, J. F. Sparks and D. Waldram, “A new infinite class of Sasaki-Einstein manifolds,” [arXiv:hep-th/0403038].
- [15] W. Chen, H. Lü, C. N. Pope and J. F. Vázquez-Poritz, “A note on Einstein-Sasaki metrics in $D \geq 7$,” Class. Quant. Grav. **22**, 3421 (2005) [arXiv:hep-th/0411218].
- [16] J. P. Gauntlett, S. Lee, T. Mateos and D. Waldram, “Marginal deformations of field theories with AdS(4) duals,” [arXiv:hep-th/0505207].
- [17] M. Cvetič, M.J. Duff, P. Hoxha, James T. Liu, H. Lü, J.X. Lu, R. Martinez-Acosta, C.N. Pope, H. Sati, Tuan A. Tran, “Embedding AdS black holes in ten and eleven dimensions,” Nucl. Phys. **B558** 96 (1999), [arXiv:hep-th/9903214].
- [18] P. Kraus, F. Larsen and S.P. Trivedi, “The Coulomb branch of gauge theory from rotating branes,” JHEP **9903** 003 (1999) [arXiv:hep-th/9811120].

- [19] M. Bershadsky, C. Vafa and V. Sadov, “D-branes and topological field theories ,” Nucl. Phys. **B463**, 420 (1996) [arXiv:hep-th/9511222].
- [20] J. Maldacena and C. Nuñez, “Supergravity description of field theories on curved manifolds and a no go theorem,” Int. J. Mod. Phys. **A16**, 822 (2001) [arXiv:hep-th/0007018].
- [21] J.B. Gutowski and H.S. Reall, “Supersymmetric AdS5 black holes,” JHEP **0402** 006 (2004) [arXiv:hep-th/0401042].
- [22] J.B. Gutowski and H.S. Reall, “General supersymmetric AdS5 black holes,” JHEP **0404** 048 (2004) [arXiv:hep-th/0401129].
- [23] J.P. Gauntlett and J.B. Gutowski, “All supersymmetric solutions of minimal gauged supergravity in five dimensions,” Phys. Rev. **D68** 105009 (2003) [arXiv:hep-th/0304064].
- [24] J.P. Gauntlett, J.B. Gutowski and N.V. Suryanarayana, “A deformation of $AdS_5 \times S^5$,” Class. Quant. Grav. **21** 5021 (2004), [arXiv:hep-th/0406188].
- [25] A.H. Chamseddine and W.A. Sabra, “Magnetic strings in five-dimensional gauged supergravity theories,” Phys. Lett. **B477**, 329 (2000) [arXiv:hep-th/9911195].
- [26] D. Klemm and W. A. Sabra, “Supersymmetry of black strings in $D = 5$ gauged supergravities,” Phys. Rev. D **62**, 024003 (2000) [arXiv:hep-th/0001131].
- [27] H. Lü, C. N. Pope and J. F. Vázquez-Poritz, “From AdS black holes to supersymmetric flux-branes,” Nucl. Phys. B **709**, 47 (2005) [arXiv:hep-th/0307001].
- [28] D. Astefanesei and G.C. Jones, “S-branes and (anti-)bubbles in (A)dS space,” JHEP **0506**, 037 (2005) [arXiv:hep-th/0502162].
- [29] H. Lin and J. Maldacena, “Fivebranes from gauge theory,” [arXiv:hep-th/0509235].
- [30] A. Khavaev and N. P. Warner, “An $N = 1$ supersymmetric Coulomb flow in IIB supergravity,” Phys. Lett. B **522**, 181 (2001) [arXiv:hep-th/0106032].
- [31] C. N. Gowdigere and N. P. Warner, “Holographic Coulomb branch flows with $N = 1$ supersymmetry,” [arXiv:hep-th/0505019].
- [32] K. Pilch and N. P. Warner, “A new supersymmetric compactification of chiral IIB supergravity,” Phys. Lett. B **487**, 22 (2000) [arXiv:hep-th/0002192].

- [33] K. Pilch and N. P. Warner, “ $N = 1$ supersymmetric renormalization group flows from IIB supergravity,” *Adv. Theor. Math. Phys.* **4**, 627 (2002) [arXiv:hep-th/0006066].
- [34] S. Frolov, “Lax pair for strings in Lunin Maldacena background,” *JHEP* **0505**, 069 (2005) [arXiv:hep-th/0503201].
- [35] S. A. Frolov, R. Roiban and A. A. Tseytlin, “Gauge - string duality for superconformal deformations of $N = 4$ super Yang-Mills theory,” [arXiv:hep-th/0503192].
- [36] S.A. Frolov, R. Roiban and A.A. Tseytlin, “Gauge string duality for (non)supersymmetric deformations of $N=4$ Super Yang-Mills theory,” [arXiv:hep-th/0507021].
- [37] M. Cvetič and D. Youm, “Rotating intersecting M-branes,” *Nucl. Phys.* **B499**, 253 (1997) [arXiv:hep-th/9612229].
- [38] M. Cvetič and S.S. Gubser, “Phases of R-charged black holes, spinning branes and strongly-coupled gauge theories,” *JHEP* **9904**, 024 (1999) [arXiv:hep-th/9902195].
- [39] K. Behrndt, A.H. Chamseddine and W.A. Sabra, “BPS black holes in $N = 2$ five-dimensional AdS supergravity,” *Phys. Lett.* **B442**, 97 (1998) [arXiv:hep-th/9807187].
- [40] D. Berenstein and R. G. Leigh, “Discrete torsion, AdS/CFT and duality,” *JHEP* **0001**, 038 (2000) [arXiv:hep-th/0001055].
- [41] D. Berenstein, V. Jejjala and R. G. Leigh, “Marginal and relevant deformations of $N = 4$ field theories and non-commutative moduli spaces of vacua,” *Nucl. Phys. B* **589**, 196 (2000) [arXiv:hep-th/0005087].
- [42] D. Berenstein, V. Jejjala and R. G. Leigh, “Noncommutative moduli spaces and T duality,” *Phys. Lett. B* **493**, 162 (2000) [arXiv:hep-th/0006168].
- [43] I. Bakas and K. Sfetsos, “States and curves of five-dimensional gauged supergravity,” *Nucl. Phys. B* **573**, 768 (2000) [arXiv:hep-th/9909041].
- [44] S. Cucu, H. Lü and J. F. Vázquez-Poritz, “Interpolating from $AdS(D-2) \times S(2)$ to $AdS(D)$,” *Nucl. Phys. B* **677**, 181 (2004) [arXiv:hep-th/0304022].
- [45] D. Z. Freedman, S. S. Gubser, K. Pilch and N. P. Warner, “Renormalization group flows from holography supersymmetry and a c-theorem,” *Adv. Theor. Math. Phys.* **3**, 363 (1999) [arXiv:hep-th/9904017].

- [46] S. Pal, “ β -deformations, potentials and KK modes,” [arXiv:hep-th/0505257].
- [47] N. Evans and J. Hockings, “ $N = 4$ Super Yang Mills at finite density: the naked truth,” JHEP **0207**, 070 (2002) [arXiv:hep-th/0205082].
- [48] H. Lin, O. Lunin and J. Maldacena, “Bubbling AdS space and 1/2 BPS geometries,” JHEP **0410**, 025 (2004) [arXiv:hep-th/0409174].
- [49] B.A. Burrington, J.T. Liu and W.A. Sabra, “ AdS_5 black holes with fermionic hair,” Phys. Rev. **D71**, 105015 (2005) [arXiv:hep-th/0412155].
- [50] S. Gukov, E. Martinec, G. Moore and A. Strominger, “The search for a holographic dual to $AdS_3 \times S^3 \times S^3 \times S^1$,” [arXiv:hep-th/0403090].
- [51] K. Pilch and N. P. Warner, “ $N = 2$ supersymmetric RG flows and the IIB dilaton,” Nucl. Phys. B **594**, 209 (2001) [arXiv:hep-th/0004063].
- [52] K. Pilch and N. P. Warner, “Generalizing the $N = 2$ supersymmetric RG flow solution of IIB supergravity,” Nucl. Phys. B **675**, 99 (2003) [arXiv:hep-th/0306098].
- [53] R. Corrado, K. Pilch and N. P. Warner, “An $N = 2$ supersymmetric membrane flow,” Nucl. Phys. B **629**, 74 (2002) [arXiv:hep-th/0107220].
- [54] C. Ahn and T. Itoh, “An $N = 1$ supersymmetric $G(2)$ -invariant flow in M-theory,” Nucl. Phys. B **627**, 45 (2002) [arXiv:hep-th/0112010].
- [55] C. Ahn and T. Itoh, “The 11-dimensional metric for AdS/CFT RG flows with common $SU(3)$ invariance,” Nucl. Phys. B **646**, 257 (2002) [arXiv:hep-th/0208137].